



A service of the National Library of Medicine  
and the National Institutes of Health

www.pubmed.gov

My NCBI  
[Sign In] [Register]

All Databases PubMed Nucleotide Protein Chemical Structure GDB PDB Journals Books

Search PubMed for Go Clear

Links Preview/Index History Clipboard Details

Display AbstractPlus Show 20 Sort by Send to

All 1 Review 0

1: J. Neuroimmunol. 2003 Dec;145(1-2):77-85.

Links

**Via beta-adrenoceptors, stimulation of extrasplenic sympathetic nerve fibers inhibits lipopolysaccharide-induced TNF secretion in perfused rat spleen.**

**Kees MG, Pongratz G, Kees F, Scholmerich J, Straub RH.**

Laboratory of Neuro/Endocrine/Immunology, Department of Internal Medicine I, University Hospital, Regensburg 93042, Germany.

Using a spleen slice microperfusion technique in mice, we have previously characterized the role of norepinephrine (NE) and other neurotransmitters for sympathetic modulation of IL-6 and TNF secretion of splenic macrophages. Since experiments in spleen slices do not reflect the situation of an entire perfused organ, we investigated sympathetic modulation of lipopolysaccharide (LPS)-induced secretion of IL-6 and TNF in perfusion experiments of rat spleen. In an organ bath, perfusion was performed in explanted whole spleens, and catecholamines and cytokines were measured by HPLC and ELISA, respectively. Release of NE depended on stimulation frequency (maximum at 10 Hz). Apart from NE, perfusates also contained significant amounts of dopamine and epinephrine. Furthermore, perfusate epinephrine levels correlated positively with perfusate NE levels (RRank=0.750,  $p<0.001$ ) but not with plasma epinephrine concentrations. This indicates that epinephrine is a conversion product of sympathetically released NE. Early electrical stimulation of extrasplenic splenic nerves, 20 min after administration of LPS, significantly inhibited TNF secretion. This electrically induced effect was abrogated by simultaneous administration of propranolol (10(-6) M) but it was not influenced by administration of either an alpha1- or alpha2-adrenergic antagonist. Late electrical stimulation of splenic nerves, 2.5 h after administration of LPS, did not change TNF secretion. Interestingly, no influence of early or late sympathetic nerve fiber stimulation on IL-6 secretion was observed. In conclusion, this is the first perfusion study of the entire spleen that demonstrates that early electrical stimulation of sympathetic splenic nerve fibers directly inhibits LPS-induced TNF secretion. This study corroborates the idea that splenic sympathetic nerves are able to inhibit important activators of the innate immune system when stimulation happens very early or even prior to their induction by LPS.

PMID: 14644033 [PubMed - indexed for MEDLINE]

Display AbstractPlus Show 20 Sort by Send to

Write to the Help Desk  
NCBI | NLM | NIH  
Department of Health & Human Services  
Privacy Statement | Freedom of Information Act | Disclaimer

#### Related Links

Neuropeptide Y cotransmission with norepinephrine in the sympathetic nerve-macrophage interplay. [J Neurocytol. 2000;]

A bacteria-induced switch of sympathetic effector mechanisms augments local inhibition of TNF-alpha and IL-6 secretion in the spleen. [FASEB J. 2000;]

Key role of the sympathetic microenvironment for the interplay of tumour necrosis factor and interleukin 6 in normal but not in inflamed mouse colon muscles. [Gut. 2005;]

Temporal profiles of interleukin-1beta, interleukin-6, and tumor necrosis factor-alpha in the plasma and hypothalamic paraventricular nucleus after intravenous or intraperitoneal administration of lipopolysaccharide in the rat: estimation by push-pull perfusion. [Endocrinology. 1999;]

Norepinephrine in mice inhibits secretion of splenic IL-6 during the dark period but stimulates its secretion in the light period--possible role of the corticosterone. [Endocrinology. 2005;]

See all Related Articles...